

BV-STRUCTURES ON THE HOMOLOGY OF THE FRAMED LONG KNOT SPACE

KEIICHI SAKAI

ABSTRACT. We introduce BV-algebra structures on the homology of the space of framed long knots in \mathbb{R}^n in two ways. The first one is given in a similar fashion to Chas-Sullivan's string topology [5]. The second one is defined on the Hochschild homology associated with a cyclic, multiplicative operad of graded modules. The latter can be applied to Bousfield-Salvatore spectral sequence converging to the homology of the space of framed long knots. Conjecturally these two structures coincide with each other.

1. INTRODUCTION

The space of framed long embeddings is known to be acted on by the little disks operad [2]. A natural question is whether this action extends to any action of the *framed* little disks operad. The answer seems affirmative, in view of [3, 14, 15], [6, 16]. In fact, Paolo Salvatore told the author that he constructed in his draft an action of the framed little disks operad on a space equivalent to the space of framed long knots, using his solution to the topological cyclic Deligne conjecture [15].

In the first result of this paper (Theorem 3.5) we imitate Chas-Sullivan's string topology [5] to realize Salvatore's homotopy-theoretical action in a geometric and homological way. Namely we define a *BV-algebra* structure on the homology of the space of framed long knots. Our BV-structure is outlined as follows. The bracket (called *Poisson bracket*) is induced by an action of little 2-disks operad [2]. The *BV-operation* (usually denoted by Δ) is derived from Hatcher's cycle [8, p. 3], which in a sense "pushes the base point through long knots". As a corollary we obtain a Lie algebra structure on the S^1 -equivariant homology. We also show that our BV-operation Δ is not trivial (Proposition 3.8 below).

Our second result (Theorem 4.6) is an algebraic one. Based on [15, 1], a homology spectral sequence, converging to the homology of the space of framed long knots (at least in higher-codimension cases), is constructed. Its E^2 -term is the *Hochschild homology* $HH_*(H_*(f\mathcal{C}))$ associated with the homology operad $H_*(f\mathcal{C})$ of the framed little disks operad, which is cyclic [3] and multiplicative. Motivated by these facts, in §4 we provide a BV-algebra structure on $HH_*(\mathcal{O})$ of any cyclic and multiplicative operad \mathcal{O} of *graded* modules. A bracket has already been defined in [17], and our BV-operation is given by a graded version of Connes' boundary operator (see [9]). Our proof is a direct analogue to that for non-graded cases [11]. Presumably Salvatore's framed little disks action would deduce the same formula as ours.

The paper is organized as follows. In §2 three spaces of framed embeddings are defined and proved to be homotopy equivalent to each other. We describe our geometric BV-algebra structure explicitly in §3; the Poisson bracket in §3.1 and

§3.2, and the Δ -operation in §3.3. The definition of $HH_*(\mathcal{O})$ is reviewed in §4.1, and the BV-algebra structure on $HH_*(\mathcal{O})$ is defined in §4.2.

2. THE SPACES OF FRAMED LONG KNOTS

We denote by $B^n := \{x \in \mathbb{R}^n \mid |x| \leq 1\}$ the n -ball, and $S^n := \partial B^{n+1} \subset \mathbb{R}^{n+1}$. We often write $\infty := (0, \dots, 0, 1) \in S^n$. S^1 is always identified with \mathbb{R}/\mathbb{Z} and let $0 = 1 \in S^1$ serve as the basepoint.

We define three spaces of framed long knots, $\widetilde{\text{Emb}}_*(S^1, S^n)$, $\widetilde{\text{Emb}}'_*(S^1, S^n)$ and $\text{EC}(1, B^{n-1})$. Eventually they turn out to be homotopy equivalent to each other. A convenient one will be used to construct each homology operation.

First we define $\text{EC}(1, B^{n-1})$, originally introduced in [2]. The homotopy type of this space can be nicely described through an action of the little disks operad, and hence its homology is equipped with a Poisson algebra structure (see §3.1).

Definition 2.1 ([2]). For a manifold M , define the space $\text{EC}(k, M)$ by

$$\text{EC}(k, M) := \{f : \mathbb{R}^k \times M \hookrightarrow \mathbb{R}^k \times M \mid f(t; x) = (t; x) \text{ if } t \notin [0, 1]^k\}.$$

We consider the case of *framed long knots*, namely $M = B^{n-1}$ and $k = 1$. The space $\text{EC}(1, B^{n-1})$ is related to the space of *long knots*

$$\mathcal{K}_n := \{f : \mathbb{R}^1 \hookrightarrow \mathbb{R}^1 \times B^{n-1} \mid f(t) = (t, 0, \dots, 0) \text{ if } t \notin [0, 1]\}$$

via the restriction map

$$\text{res} : \text{EC}(1, B^{n-1}) \rightarrow \mathcal{K}_n, \quad f \mapsto f|_{\mathbb{R}^1 \times \{0\}}$$

which is a fibration with fiber $\Omega SO(n-1)$.

Definition 2.2. Define $\iota : S^1 \hookrightarrow S^n$ and $\tilde{\iota} : S^1 \times B^{n-1} \hookrightarrow S^n$ by

$$\begin{aligned} \iota(t) &:= (\sin 2\pi t, 0, \dots, 0, \cos 2\pi t) \in S^n \subset \mathbb{R}^{n+1}, \quad \text{and} \\ \tilde{\iota}(t; x) &:= \frac{\iota(t) + \epsilon_0(0, x, 0)}{|\iota(t) + \epsilon_0(0, x, 0)|}, \end{aligned}$$

where $\epsilon_0 > 0$ is some fixed small number. Define $\widetilde{\text{Emb}}'_*(S^1, S^n)$ to be the space of embeddings $\tilde{\varphi} : S^1 \times B^{n-1} \hookrightarrow S^n$ such that $\tilde{\varphi}(0; x) = \tilde{\iota}(0; x)$ and all the partial derivatives of $\tilde{\varphi}$ at $(0; x) \in S^1 \times B^{n-1}$ of all orders are equal to those of $\tilde{\iota}$. Similarly define $\text{Emb}'_*(S^1, S^n)$ to be the space of all embeddings $\varphi : S^1 \hookrightarrow S^n$ such that $\varphi(0) = \iota(0) = \infty$ and all of its derivatives at $t = 0$ are equal to those of ι .

The restriction map

$$\text{res} : \widetilde{\text{Emb}}'_*(S^1, S^n) \rightarrow \text{Emb}'_*(S^1, S^n), \quad \tilde{\varphi} \mapsto \tilde{\varphi}|_{S^1 \times \{0\}},$$

is also a fibration with fiber $\Omega SO(n-1)$.

There is another embedding space on which S^1 acts (Lemma 3.4):

Definition 2.3. Define $\text{Emb}_*(S^1, S^n)$ to be the space of embeddings $\varphi : S^1 \hookrightarrow S^n$ satisfying

$$\varphi(0) = \infty, \quad \varphi'(0)/|\varphi'(0)| = (1, 0, \dots, 0).$$

Define $\widetilde{\text{Emb}}_*(S^1, S^n)$ to be the space of pairs $(\varphi; w)$, where $\varphi \in \text{Emb}_*(S^1, S^n)$ and $w = (w_0, \dots, w_n) : S^1 \rightarrow SO(n+1)$ ($w_i : S^1 \rightarrow S^n$) satisfying

- $w_0(t) = \varphi'(t)/|\varphi'(t)|$ and $w_n(t) = \varphi(t)$ for any $t \in S^1$,
- $w(0) = I_{n+1}$ (the identity matrix).

Denote by $\pi : \widetilde{\text{Emb}}_*(S^1, S^n) \rightarrow \text{Emb}_*(S^1, S^n)$ the natural projection.

Define $\tilde{\text{cl}} : \text{EC}(1, B^{n-1}) \rightarrow \widetilde{\text{Emb}}'_*(S^1, S^n)$ (“closure” of long embeddings) by $\tilde{\text{cl}}(f) := \tilde{\iota} \circ f$. This is smooth at $t = 0$ and has the same partial derivatives at $(0; x) \in S^1 \times B^{n-1}$ as $\tilde{\iota}$ because $f \in \text{EC}(1, B^{n-1})$ extends to $\text{id}_{\mathbb{R}^1 \times B^{n-1}}$ outside $[0, 1]$. Similarly define the map $\text{cl} : \mathcal{K}_n \rightarrow \text{Emb}'_*(S^1, S^n)$ by $\text{cl}(f) := \tilde{\iota} \circ f$.

Define $\tilde{h} : \widetilde{\text{Emb}}'_*(S^1, S^n) \rightarrow \widetilde{\text{Emb}}_*(S^1, S^n)$ by

$$\tilde{h}(\tilde{\varphi})(t) := \left(\tilde{\varphi}(t; 0); GS \left(\frac{\partial \tilde{\varphi}}{\partial t}(t; 0), \frac{\partial \tilde{\varphi}}{\partial x_1}(t; 0), \dots, \frac{\partial \tilde{\varphi}}{\partial x_{n-1}}(t; 0), \tilde{\varphi}(t; 0) \right) \right),$$

where $GS : GL_{n+1}^+(\mathbb{R}) \xrightarrow{\cong} SO(n+1)$ is the Gram-Schmidt orthonormalization, and define $h : \text{Emb}'_*(S^1, S^n) \rightarrow \text{Emb}_*(S^1, S^n)$ as the natural inclusion. Note that we have maps of fibration sequences;

$$(2.1) \quad \begin{array}{ccccc} \Omega SO(n-1) & \longrightarrow & \text{EC}(1, B^{n-1}) & \xrightarrow{\text{res}} & \mathcal{K}_n \\ \parallel & & \tilde{\text{cl}} \downarrow & & \text{cl} \downarrow \\ \Omega SO(n-1) & \longrightarrow & \widetilde{\text{Emb}}'_*(S^1, S^n) & \xrightarrow{\text{res}} & \text{Emb}'_*(S^1, S^n) \\ \parallel & & \tilde{h} \downarrow & & h \downarrow \\ \Omega SO(n-1) & \longrightarrow & \widetilde{\text{Emb}}_*(S^1, S^n) & \xrightarrow{\pi} & \text{Emb}_*(S^1, S^n) \end{array}$$

Proposition 2.4. *The maps $\tilde{\text{cl}}$ and cl are homotopy equivalences.*

Proof. Because the embedding spaces have homotopy types of CW-complexes, it suffices to show that the maps $\tilde{\text{cl}}$ and cl are weak homotopy equivalences. By (2.1), it is enough to show that cl is a (weak) homotopy equivalence.

The homotopy inverse cl^{-1} is given by

$$\text{cl}(\varphi)(t) := \begin{cases} \Phi(\varphi(t)) & \text{if } t \in [0, 1], \\ (t, 0, \dots, 0) & \text{if } t \notin [0, 1], \end{cases}$$


where $\Phi : S^n \setminus \{\infty\} \xrightarrow{\cong} (0, 1) \times \text{Int } B^{n-1}$ is the stereographic projection $S^n \setminus \{\infty\} \xrightarrow{\cong} \mathbb{R}^n$ followed by a dilation $\mathbb{R}^n \xrightarrow{\cong} (0, 1) \times \text{Int } B^{n-1}$ which is chosen so that $\Phi(\iota(t)) = (t, 0, \dots, 0)$ for $t \in [0, 1]$. \square

Proposition 2.5. *The maps \tilde{h} and h are homotopy equivalences.*

Proof. Similarly to the proof of Proposition 2.4, it is enough to show that h is a weak homotopy equivalence. The idea is to “straighten” the elements of $\text{Emb}_*(S^1, S^n)$ around $t = 0$.

First we show that $h_* : \pi_k(\text{Emb}'_*(S^1, S^n), \iota) \rightarrow \pi_k(\text{Emb}_*(S^1, S^n), \iota)$ is surjective (the basepoint ι will be omitted below). Let $\xi \in \pi_k(\text{Emb}_*(S^1, S^n))$ be represented by $g : S^k \rightarrow \text{Emb}_*(S^1, S^n)$. For each $z \in S^k$, there exists $\epsilon > 0$ such that the ϵ -ball $B_\epsilon \subset S^n$ (with respect to the standard metric on S^n) centered at $\infty \in S^n$ satisfies that

- $g(z)(S^1) \cap B_\epsilon$ is connected, and
- if $g(z)^{-1}(B_\epsilon) = (-\delta_1, \delta_2)$, then $g(z)_1$ (the first coordinate of $g(z)$) monotonically increases on the interval $(-\delta_1, \delta_2)$.

FIGURE 2.1. A modification of g to be standard near ∞

Such an $\epsilon > 0$ as above can be taken uniformly for all $z \in S^k$. Indeed there is $\epsilon' > 0$ such that $g(z)'_1(t) > 0$ for all $(z, t) \in S^k \times (-\epsilon', \epsilon')$, because the map $S^k \times S^1 \rightarrow \mathbb{R}$ given by $(z, t) \mapsto g(z)'_1(t)$ is continuous and $\{g(z)'_1(0) \mid z \in S^k\}$ has the positive minimum by the compactness of S^k . Then we can take $\epsilon > 0$ so that B_ϵ does not intersect the compact set $\hat{g}(S^k \times (S^1 \setminus (-\epsilon', \epsilon')))$, where $\hat{g}(z, t) := g(z)(t)$. This ϵ satisfies the above conditions. Note that $\delta_1, \delta_2 > 0$ depend continuously on z .

Consider a natural projection $p : B_\epsilon \rightarrow \iota(S^1) \cap B_\epsilon$ and diffeomorphisms $s_z : (-\delta_1, \delta_2) \rightarrow (-\delta'_1, \delta'_2)$ for some $\delta'_1, \delta'_2 > 0$ such that $p(g(z)(t)) = \iota(s_z(t))$ (see Figure 2.1). Putting $\delta := \min_{z \in S^k} \{\delta_1, \delta_2\}$, define

$$\bar{g}(z)(t) := \frac{(1 - b_\delta(t))g(z)(t) + b_\delta(t)\iota(s_z(t))}{|(1 - b_\delta(t))g(z)(t) + b_\delta(t)\iota(s_z(t))|}$$

where $b_\delta(t) := b(t/\delta)$ and $b : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ is a fixed bump function with support $|t| \leq 1$ satisfying $b(t) \equiv 1$ for $|t| \leq 1/2$ (and hence $b_\delta(t) \equiv 1$ for $|t| \leq \delta/2$). By construction \bar{g} is homotopic to g and $\bar{g}(z)(t) = \iota(s_z(t))$ on $(-\delta, \delta)$. Define

$$\tilde{g}(z)(t) := \bar{g}(z)(b_{\delta/2}(t)t + (1 - b_{\delta/2}(t))s_z(t))$$

then $g \sim \tilde{g}$ and $\tilde{g}(z) = \iota$ on $(-\delta/4, \delta/4)$, and hence \tilde{g} represents an element of $\pi_k(\text{Emb}'_*(S^1, S^n))$ which is mapped to $\xi = [g]$. Thus surjectivity follows.

Injectivity is proved in a similar way. Suppose $\eta = [g] \in \pi_k(\text{Emb}'_*(S^1, S^n))$ maps to $0 \in \pi_k(\text{Emb}_*(S^1, S^n))$, and choose a map $G : B^{k+1} \rightarrow \text{Emb}_*(S^1, S^n)$ bounded by g . Since B^{k+1} is compact, we can deform G to be standard near $t = 0$ in a similar way to the above. Thus g bounds a ball in $\text{Emb}'_*(S^1, S^n)$ and hence $\eta = 0$. \square

Remark 2.6. In fact the homotopy inverse $\tilde{\text{cl}}^{-1}$ is given by composing an appropriate diffeomorphism $\Psi : S^n \setminus \tilde{\iota}(\{0\} \times B^{n-1}) \xrightarrow{\cong} (0, 1) \times \text{Int } B^{n-1}$ to the elements of $\widetilde{\text{Emb}}'_*(S^1, S^n)$. The homotopy inverse \tilde{h}^{-1} is given by “straightening” the elements of $\widetilde{\text{Emb}}_*(S^1, S^n)$ and “fattening” the knots by geodesics along the framings.


3. A GEOMETRIC BV-STRUCTURE

Definition 3.1. A k -Poisson (-Gerstenhaber) algebra A is a graded commutative algebra equipped with a graded Lie bracket $[-, -] : A \times A \rightarrow A$ of degree k , called Poisson (-Gerstenhaber) bracket, satisfying the Leibniz rule

$$[x, yz] = [x, y]z + (-1)^{(\tilde{x}+k)\tilde{y}}y[x, z],$$

where \tilde{x} denotes the degree of x , that is, $x \in A_{\tilde{x}}$. A 1-Poisson algebra A is called a BV-algebra if it is endowed with a degree one operation $\Delta : A \rightarrow A$ satisfying

- $\Delta \circ \Delta = 0$,
- $\Delta(xy) = \Delta(x)y + (-1)^{\tilde{x}}x\Delta(y) + (-1)^{\tilde{x}}[x, y]$.


 FIGURE 3.1. Connected-sums on $\text{EC}(k, M)$ and $\widetilde{\text{Emb}}_*(S^1, S^n)$

The aim of this section is to define a BV-algebra structure on $H_*(\widetilde{\text{Emb}}_*(S^1, S^n))$. A Poisson algebra structure on $H_*(\text{EC}(1, B^{n-1}))$ has already been defined in [3]. First we describe this structure on $H_*(\widetilde{\text{Emb}}'_*(S^1, S^n))$ (§3.2). Then we define the Δ -operation using an S^1 -action on $\widetilde{\text{Emb}}_*(S^1, S^n)$.

3.1. Budney's Poisson structure. Budney [2] constructed an action of the little $(k+1)$ -disks operad \mathcal{C}_{k+1} on $\text{EC}(k, M)$. The main idea of the action can be found in [2, Fig. 2]; start with the *connected-sum* $f\sharp g$ (defined explicitly below; see Figure 3.1), “push off” g through f as in the right-half of [2, Fig. 2] until we arrive at $g\sharp f$, and perform the same procedure with f and g exchanged. As a corollary we have the following.

Theorem 3.2 ([2]). *$H_*(\text{EC}(k, M))$ admits a k -Poisson algebra structure.*

Here we describe Budney's Poisson algebra structure on $H_*(\text{EC}(1, B^{n-1}))$ explicitly. The product (denoted by $x \cdot y$, or simply xy) and the Poisson bracket (denoted by λ) are induced by the “second stage” of the action of \mathcal{C}_2 ;

$$\mu_2 : \mathcal{C}_2(2) \times \text{EC}(1, B^{n-1})^{\times 2} \rightarrow \text{EC}(1, B^{n-1}).$$

The product corresponds to the generator of $H_0(\mathcal{C}_2(2)) \cong \mathbb{Z}$ and is induced by the *connected-sum* defined as follows. For $0 \leq \alpha \leq 1$, define two diffeomorphisms

$$\begin{aligned} l_\alpha : \mathbb{R}^1 &\xrightarrow{\cong} \mathbb{R}^1 && \text{by } l_\alpha(t) := 2t - \alpha, \\ s_\alpha : \mathbb{R}^1 \times B^{n-1} &\xrightarrow{\cong} \mathbb{R}^1 \times B^{n-1} && \text{by } s_\alpha := l_\alpha \times \text{id}_{B^{n-1}}. \end{aligned}$$

Then for $f \in \text{EC}(1, B^{n-1})$,

$$L_\alpha f := s_\alpha^{-1} \circ f \circ s_\alpha \in \text{EC}(1, B^{n-1})$$

has the support $[\frac{\alpha}{2}, \frac{\alpha+1}{2}] \times B^{n-1}$ and satisfies

$$L_\alpha f([\frac{\alpha}{2}, \frac{\alpha+1}{2}] \times B^{n-1}) \subset [\frac{\alpha}{2}, \frac{\alpha+1}{2}] \times B^{n-1}$$

(see [2, Fig. 4]). Then for $f, g \in \text{EC}(1, B^{n-1})$, define $f\sharp g \in \text{EC}(1, B^{n-1})$ by

$$f\sharp g(t; x) := \begin{cases} L_0 f(t; x) & t \leq \frac{1}{2}, \\ L_1 g(t; x) & t \geq \frac{1}{2} \end{cases}$$

(see Figure 3.1). We notice that the elements in $\text{EC}(1, B^{n-1})$ are maps $\mathbb{R}^1 \times B^{n-1} \rightarrow \mathbb{R}^1 \times B^{n-1}$ and we can compose them. Using the composition, we can write $f\sharp g$ as

$$f\sharp g = (L_0 f) \circ (L_1 g) = (L_1 g) \circ (L_0 f).$$

Poisson bracket λ corresponds to the generator of $H_1(\mathcal{C}_2(2)) \cong \mathbb{Z}$ and can be described as follows. For $0 \leq \alpha \leq 1$ and $f, g \in \text{EC}(1, B^{n-1})$, define $f *_\alpha g \in \text{EC}(1, B^{n-1})$ by

$$f *_\alpha g := (L_\alpha f) \circ (L_{1-\alpha} g).$$

This defines the **-operation*

$$* : I \times \text{EC}(1, B^{n-1})^{\times 2} \rightarrow \text{EC}(1, B^{n-1}), \quad (\alpha, f, g) \mapsto f *_\alpha g.$$

Remark 3.3. The **-operation* gives a homotopy between $f *_0 g = f \sharp g$ and $f *_1 g = g \sharp f$, and hence \sharp is homotopy commutative. In particular for homology classes x, y , we have $x *_\alpha y = xy$ for any $0 \leq \alpha \leq 1$.

The map $S^1 \times \text{EC}(1, B^{n-1})^{\times 2} \rightarrow \text{EC}(1, B^{n-1})$ defined by

$$(\alpha, f, g) \mapsto \begin{cases} f *_\alpha g & 0 \leq \alpha \leq \frac{1}{2}, \\ g *_\alpha f & \frac{1}{2} \leq \alpha \leq 1 \end{cases}$$

corresponds to the map μ_2 appearing in Budney's \mathcal{C}_2 -action, and the Poisson bracket λ is induced by this map and a fixed generator of $H_1(S^1) \cong \mathbb{Z}$.

For any cubical chains $\xi : I^p \rightarrow \text{EC}(1, B^{n-1})$ and $\eta : I^q \rightarrow \text{EC}(1, B^{n-1})$, define cubical $(p+q+1)$ -chains $\xi * \eta$ and $\eta *' \xi$ by

$$\begin{aligned} \xi * \eta &: I \times I^p \times I^q \ni (\alpha, u, v) \mapsto \xi(u) *_\alpha \eta(v), \\ \eta *' \xi &: I \times I^p \times I^q \ni (\alpha, u, v) \mapsto \eta(v) *_\alpha \xi(u), \end{aligned}$$

and extend them linearly on the cubical chain complex. We also define cubical $(p+q)$ -chains $\xi *_\alpha \eta := \xi * \eta|_{\{\alpha\} \times I^{p+q}}$ and $\eta *_\alpha' \xi := \eta *' \xi|_{\{\alpha\} \times I^{p+q}}$. If x and y are cubical p - and q -cycles, then $\partial(x * y) = x *_1 y - x *_0 y$ and $\partial(y *' x) = y *_1' x - y *_0' x$. Since $y *_1' x = x *_0 y$ and $y *_0' x = x *_1 y$,

$$(3.1) \quad (-1)^{p-1} (x * y + y *' x)$$

is a cubical $(p+q+1)$ -cycle. The cycle (3.1) represents $\lambda(x, y)$.

3.2. Poisson structure for $\widetilde{\text{Emb}}'_*(S^1, S^n)$. We can transfer the above Poisson structure on $H_*(\text{EC}(1, B^{n-1}))$ to $H_*(\widetilde{\text{Emb}}'_*(S^1, S^n))$ via the homotopy equivalence $\tilde{\text{cl}}$ defined in Proposition 2.4. We illustrate this structure from a geometric view.

Figure 3.1 explains the idea of connected-sum on $\widetilde{\text{Emb}}'_*(S^1, S^n)$. For $\tilde{\varphi} \in \widetilde{\text{Emb}}'_*(S^1, S^n)$ and $0 \leq \alpha \leq 1$, define

$$L_\alpha \tilde{\varphi} := \tilde{\text{cl}}(L_\alpha \tilde{\text{cl}}^{-1}(\tilde{\varphi})).$$

Roughly speaking $L_\alpha \tilde{\varphi}$ is $\tilde{\varphi}$ with the cylinder $\tilde{\iota}([\frac{\alpha}{2}, \frac{\alpha+1}{2}] \times B^{n-1}) \cong I \times B^{n-1}$ replaced by $\tilde{\text{cl}}^{-1}(\tilde{\varphi})$. Then for $\tilde{\varphi}_1, \tilde{\varphi}_2 \in \widetilde{\text{Emb}}'_*(S^1, S^n)$,

$$(3.2) \quad \tilde{\varphi}_1 \sharp \tilde{\varphi}_2 := (L_0 \tilde{\varphi}_1) \circ (L_1 \tilde{\varphi}_2) = \begin{cases} L_0 \tilde{\varphi}_1 & \text{on } [0, \frac{1}{2}] \times B^{n-1}, \\ L_1 \tilde{\varphi}_2 & \text{on } [\frac{1}{2}, 1] \times B^{n-1}. \end{cases}$$

The **-operation* for $\widetilde{\text{Emb}}'_*(S^1, S^n)$

$$* : I \times \widetilde{\text{Emb}}'_*(S^1, S^n)^{\times 2} \rightarrow \widetilde{\text{Emb}}'_*(S^1, S^n)$$

is defined by using that for $\text{EC}(1, B^{n-1})$;

$$(3.3) \quad \begin{aligned} (\alpha, \tilde{\varphi}_1, \tilde{\varphi}_2) &\mapsto \tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2 := \tilde{\text{cl}}(\tilde{\text{cl}}^{-1}(\tilde{\varphi}_1) *_\alpha \tilde{\text{cl}}^{-1}(\tilde{\varphi}_2)) \\ &= (L_\alpha \tilde{\varphi}_1) \circ (L_{1-\alpha} \tilde{\text{cl}}^{-1}(\tilde{\varphi}_2)). \end{aligned}$$

Roughly speaking $\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2$ is $L_\alpha \tilde{\varphi}_1$ with $L_\alpha \tilde{\varphi}_1([\frac{1-\alpha}{2}, \frac{2-\alpha}{2}] \times B^{n-1})$ replaced by $\text{cl}^{-1}(\tilde{\varphi}_2)$.

3.3. BV-operation. We define an S^1 -action on $\widetilde{\text{Emb}}_*(S^1, S^n)$ as was done in [8, p. 3]. This action induces our BV-operation on $H_*(\widetilde{\text{Emb}}_*(S^1, S^n))$.

For any $(\varphi; w) \in \widetilde{\text{Emb}}_*(S^1, S^n)$ and $\alpha \in S^1$, define $(\varphi; w)^\alpha \in \widetilde{\text{Emb}}_*(S^1, S^n)$ by

$$(\varphi; w)^\alpha(t) := (A\varphi(t - \alpha); Aw(t - \alpha)),$$

where $A = w(-\alpha)^{-1} \in SO(n+1)$ (acting on S^n in the usual way). Since $Aw(-\alpha) = I_{n+1}$ (and hence $A\varphi(-\alpha) = \infty$), $(\varphi; w)^\alpha$ is indeed in $\widetilde{\text{Emb}}_*(S^1, S^n)$.

Lemma 3.4. *The above formula defines an S^1 -action on $\widetilde{\text{Emb}}_*(S^1, S^n)$. That is, we have $((\varphi; w)^\alpha)^\beta = (\varphi; w)^{\alpha+\beta}$ and $(\varphi; w)^0 = (\varphi; w)$.*

This action can be interpreted for $\tilde{\varphi} \in \widetilde{\text{Emb}}'_*(S^1, S^n)$ via \tilde{h} (see (2.1));

$$\tilde{\varphi}^\alpha := \tilde{h}^{-1}(\tilde{h}(\tilde{\varphi})^\alpha).$$

The fundamental class of S^1 induces our Δ -operation through the above action;

$$\Delta : H_*(\widetilde{\text{Emb}}_*(S^1, S^n)) \rightarrow H_{*+1}(\widetilde{\text{Emb}}_*(S^1, S^n)).$$

We have $\Delta^2 = 0$ since Δ is induced by an S^1 -action and $H_*(S^1) = \bigwedge^* \langle [S^1] \rangle$.

Theorem 3.5. *$(H_*(\widetilde{\text{Emb}}_*(S^1, S^n)), \cdot, \lambda, \Delta)$ is a BV-algebra.*

Proof. We need to prove the last equality of Definition 3.1, that is, Δ is a derivation with respect to the product modulo λ ;

$$(3.4) \quad \Delta(xy) - \Delta(x)y - (-1)^{\bar{x}}x\Delta(y) = (-1)^{\bar{x}}\lambda(x, y).$$

This is proved in a similar way to [5, Lemma 5.2]. Define two operations Δ_i ($i = 1, 2$) as the “first/last half” of Δ ;

$$\Delta_i : I \times \widetilde{\text{Emb}}_*(S^1, S^n) \rightarrow \widetilde{\text{Emb}}_*(S^1, S^n), \quad \Delta_i(\alpha, \sigma) = \sigma^{(\alpha+i-1)/2}.$$

Let $\Delta^2 := \{0 \leq \beta \leq \alpha \leq 1\}$ be the standard 2-simplex. Define

$$\Phi : \Delta^2 \times \widetilde{\text{Emb}}_*(S^1, S^n)^{\times 2} \rightarrow \widetilde{\text{Emb}}_*(S^1, S^n)$$

by

$$\Phi((\alpha, \beta), \sigma_1, \sigma_2) := (\sigma_1 *_\alpha \sigma_2)^{\beta/2}.$$

Choose cubical p - and q -cycles $x = \sum a_i x_i$ and $y = \sum b_j y_j$. Then $\Phi_*(\text{id}, x, y) = \sum a_i b_j \Phi \circ (\text{id}_{\Delta^2} \times x_i \times y_j)$ is a $(p+q+2)$ -chain whose boundary comes from $\partial \Delta^2$. We can see that

- (i) $\{\alpha = \beta\} \subset \partial \Delta^2$ corresponds to a $(p+q+1)$ -cycle homologous to $\Delta(x)y$,
- (ii) $\{\alpha = 1\} \cup \{\beta = 0\} \subset \partial \Delta^2$ corresponds to a $(p+q+1)$ -cycle homologous to $x * y + \Delta_1(x *_1 y)$ (see Figure 3.2).

(ii) is immediate from the definition, and (i) follows from Lemma 3.6 below. Thus

$$(3.5) \quad x * y + \Delta_1(x *_1 y) - \Delta(x)y \sim 0.$$

Similarly, considering the $(p+q+2)$ -chain $(y *_\alpha' x)^{\beta/2}$, we have

$$(3.6) \quad y *_1' x + \Delta_1(y *_1' x) - \Delta(y) *_0' x \sim 0.$$

By definition we have equalities of $(p+q+1)$ -chains

$$(3.7) \quad \Delta_1(y *_1' x) = \Delta_2(y *_0' x) = \Delta_2(x *_1 y).$$




FIGURE 3.2. The chain $\Phi_*(\text{id}, x, y)$; the symbol \times indicates the basepoint

We also see, by exchanging $u \in I^p$ and $v \in I^q$, that the cycle $\Delta(y) *'_0 x$ is homologous to $(-1)^{pq} \Delta(y) *'_0 x \sim (-1)^p x \Delta(y)$. Substituting this and (3.7) into (3.6) we have

$$(3.8) \quad y *'_0 x + \Delta_2(x *_1 y) - (-1)^p x \Delta(y) \sim 0.$$

Then (3.5), (3.8) and the facts $\Delta_1(z) + \Delta_2(z) \sim \Delta(z)$ (for any cycle z) and $x *_1 y \sim xy$ (see Remark 3.3) imply (3.4). \square

Lemma 3.6. $\Phi|_{\{\alpha=\beta\} \times \widetilde{\text{Emb}}_*(S^1, S^n)^{\times 2}} : I \times \widetilde{\text{Emb}}'_*(S^1, S^n)^{\times 2} \rightarrow \widetilde{\text{Emb}}'_*(S^1, S^n)$, defined by $(\alpha, \tilde{\varphi}_1, \tilde{\varphi}_2) \mapsto (\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2)^{\alpha/2}$, is homotopic to $(\alpha, \tilde{\varphi}_1, \tilde{\varphi}_2) \mapsto (\tilde{\varphi}_1)^\alpha \# \tilde{\varphi}_2$.

Proof. For $\tilde{\varphi}_1, \tilde{\varphi}_2 \in \widetilde{\text{Emb}}'_*(S^1, S^n)$, the embedding $\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2$ is given by (3.3), which is equal to $L_\alpha \tilde{\varphi}_1$ at $t = -\alpha/2 \equiv \frac{2-\alpha}{2}$. The embedding $\tilde{h}(\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2)^{\alpha/2} \in \widetilde{\text{Emb}}'_*(S^1, S^n)$ is $\tilde{h}(\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2)$ rotated by an action of a matrix which is determined by the value $\tilde{h}(\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2)(-\alpha/2) = \tilde{h}(L_\alpha \tilde{\varphi}_1)(-\alpha/2)$. Thus $(\tilde{\varphi}_1 *_\alpha \tilde{\varphi}_2)^{\alpha/2}$ is homotopic to

$$\begin{aligned} (t; x) &\mapsto A(L_\alpha \tilde{\varphi}_1(-\alpha/2)) \cdot (L_\alpha \tilde{\varphi}_1(L_{1-\alpha} \tilde{\text{cl}}^{-1}(\tilde{\varphi}_2)(t - \frac{\alpha}{2}, x))) \\ &= (L_\alpha \tilde{\varphi}_1)^{\alpha/2} (L_1 \tilde{\text{cl}}^{-1}(\tilde{\varphi}_2)(t, x)) \\ &\sim (L_0 \tilde{\varphi}_1)^\alpha (L_1 \tilde{\text{cl}}^{-1}(\tilde{\varphi}_2)(t, x)), \end{aligned}$$

here the equality follows from $l_{1-\alpha}(t - \frac{\alpha}{2}) = l_1(t)$ and $l_{1-\alpha}^{-1}(u) = l_1^{-1}(u) - \frac{\alpha}{2}$, and then we use Lemma 3.7 below. The proof is completed by considering the homotopy from any $f \in \text{EC}(1, B^{n-1})$ to $L_0 f$ given by

$$\beta \mapsto \hat{s}_\beta^{-1} \circ f \circ \hat{s}_\beta$$

where $\hat{s}_\beta := \hat{l}_\beta \times \text{id}_{B^{n-1}}$ and $\hat{l}_\beta(t) := (1 + \beta)t$, and translating this homotopy to a homotopy on $\widetilde{\text{Emb}}'_*(S^1, S^n)$ via $\tilde{\text{cl}}$. \square

Lemma 3.7. The map $I \times \widetilde{\text{Emb}}'_*(S^1, S^n) \rightarrow \widetilde{\text{Emb}}'_*(S^1, S^n)$ given by $(\alpha, \sigma) \mapsto (L_\alpha \sigma)^{\alpha/2}$ (where $L_\alpha \sigma$ is defined through the homotopy equivalence \tilde{h}) is equal to $(\alpha, \sigma) \mapsto (L_0 \sigma)^\alpha$.

Proof. First by definition $L_\alpha\sigma(t) = R_\alpha \cdot L_0\sigma(t - \frac{\alpha}{2})$ where R_α is the rotation by $\pi\alpha$ in the x_1x_{n+1} -plane. Putting $L_0\sigma = (\varphi; w)$, we have

$$(L_\alpha\sigma)^{\alpha/2}(t) = (AR_\alpha\varphi((t - \frac{\alpha}{2}) - \frac{\alpha}{2}); AR_\alpha w((t - \frac{\alpha}{2}) - \frac{\alpha}{2}))$$

where $A = (R_\alpha w(-\frac{\alpha}{2} - \frac{\alpha}{2}))^{-1} = w(-\alpha)^{-1}R_\alpha^{-1}$. Thus

$$(L_\alpha\sigma)^{\alpha/2}(t) = (w(-\alpha)^{-1}\varphi(t - \alpha); w(-\alpha)^{-1}w(t - \alpha)) = (L_0\sigma)^\alpha(t). \quad \square$$

Proposition 3.8. Δ is nontrivial when $n \geq 3$ is odd.

Proof. It is an easy consequence of the third equation from Definition 3.1 that, if $\lambda(x, y) \neq 0$, then one of $\Delta(xy)$, $\Delta(x)$ and $\Delta(y)$ is not zero. The non-triviality of λ is proved in [4] (when $n = 3$) and in [13] (when $n > 3$ is odd). \square

3.4. The string bracket. Similarly to [5, §6], consider the principal S^1 -bundle

$$\pi : ES^1 \times \widetilde{\text{Emb}}_*(S^1, S^n) \rightarrow ES^1 \times_{S^1} \widetilde{\text{Emb}}_*(S^1, S^n).$$

Let $p : E \rightarrow ES^1 \times_{S^1} \widetilde{\text{Emb}}_*(S^1, S^n)$ be the vector bundle of rank two associated with π , and E_0 the complement of the zero section of E . The Gysin exact sequence for p can be written as

$$\begin{aligned} \dots \rightarrow H_i(\widetilde{\text{Emb}}_*(S^1, S^n)) \xrightarrow{E} H_i^{S^1}(\widetilde{\text{Emb}}_*(S^1, S^n)) \\ \xrightarrow{c} H_{i-2}^{S^1}(\widetilde{\text{Emb}}_*(S^1, S^n)) \xrightarrow{M} H_{i-1}(\widetilde{\text{Emb}}_*(S^1, S^n)) \rightarrow \dots \end{aligned}$$

(E is induced by $E_0 \hookrightarrow E$, c is given by capping the Euler class, and M is the connecting homomorphism). Define the bracket $\{-, -\}$ on $H_*^{S^1}(\widetilde{\text{Emb}}_*(S^1, S^n))$ by

$$\{x, y\} := (-1)^{\tilde{x}} E(M(x)M(y)).$$

As a corollary of Theorem 3.5 we obtain the following.

Corollary 3.9. $\{-, -\}$ is a degree two Lie bracket on $H_*^{S^1}(\widetilde{\text{Emb}}_*(S^1, S^n))$.

Proof. The proof is the same as that of [5, Theorem 6.1], which formally uses the BV-structure on $H_*(\widetilde{\text{Emb}}_*(S^1, S^n))$, the fact that the map Δ induced by the S^1 -action can be described as $\Delta = M \circ E$, and the exactness of the Gysin sequence. \square

4. BV-STRUCTURE ON THE HOCHSCHILD HOMOLOGY

This section is independent of the previous one. In this section we define a BV-algebra structure on the Hochschild homology $HH_*(\mathcal{O})$ associated with a cyclic multiplicative operad \mathcal{O} in the category of graded modules.

One motivation is as follows. When $n \geq 4$, the space $\text{EC}(1, B^{n-1})$ is weakly equivalent to the homotopy totalization of an operad fK_n , called the *framed Kontsevich operad*, which is weakly equivalent to the framed little n -disks operad $f\mathcal{C}_n$ [14]. There is a spectral sequence [1] converging to the homology of the homotopy totalization of a topological multiplicative operad (fK_n is one of such operads), and its E^2 -term is the Hochschild homology associated with the homology of the operad. In general, for any multiplicative operad \mathcal{O} of modules, its Hochschild homology $HH_*(\mathcal{O})$ admits a Poisson algebra structure [17], and if moreover \mathcal{O} is a cyclic operad over *non-graded* modules, then $HH_*(\mathcal{O})$ admits a BV-algebra structure [18, 11]. For $\text{EC}(1, B^{n-1})$, the operad $\mathcal{O} = H_*(f\mathcal{C}_n)$ is a multiplicative operad of graded modules, and the Poisson structure on $HH_*(\mathcal{O})$ is proved in Salvatore's

draft to coincide with that described in [2]. Moreover, $f\mathcal{C}_n$ is equivalent to a cyclic operad (of “conformal n -balls”) [3], and it turns out that $H_*(f\mathcal{C}_n)$ is a cyclic multiplicative operad of *graded* modules. So it is natural to ask whether $HH_*(\mathcal{O})$ admits a suitable BV-algebra structure when \mathcal{O} is a cyclic operad of *graded* modules, in such a way that it coincides with that discussed in §3 in the case of embedding spaces. Our construction is a direct analogue to the non-graded cases.

As for operads, we follow the convention of [10].

4.1. Hochschild homology. For an operad \mathcal{O} and $x \in \mathcal{O}(l)$, $y \in \mathcal{O}(m)$, define

$$x \circ_i y := x(\text{id}, \dots, \text{id}, y, \text{id}, \dots, \text{id}) \in \mathcal{O}(l+m-1),$$

where y sits in the i -th place, and $\text{id} \in \mathcal{O}(1)$ is the identity element. When \mathcal{O} is an operad of graded modules, we denote by \tilde{x} the grading of x in the graded module $\mathcal{O}(l)$, that is, $x \in \mathcal{O}(l)_{\tilde{x}}$.

Let \mathcal{O} be a multiplicative operad [10, Definition 10.1] of graded modules; namely \mathcal{O} is a non-symmetric operad of graded modules endowed with a morphism $\mathcal{ASSOC} \rightarrow \mathcal{O}$, where \mathcal{ASSOC} is the *associative operad* given by $\mathcal{ASSOC}(n) = \{*\}$ for all $n \geq 0$. We denote the image of $\mathcal{ASSOC}(2) = \{*\} \rightarrow \mathcal{O}(2)$ by $\mu \in \mathcal{O}(2)$ and call it the *multiplication*. The collection $\mathcal{O} = \{\mathcal{O}(k)\}_{k \geq 0}$ admits a cosimplicial module structure; the cosimplicial structure maps

$$d^i : \mathcal{O}(k-1) \rightarrow \mathcal{O}(k), \quad s^i : \mathcal{O}(k+1) \rightarrow \mathcal{O}(k) \quad (0 \leq i \leq k)$$

are defined as in [10, §10] by using μ and the unit element $e \in \mathcal{O}(0)$. The grading-preserving map

$$\partial_k : \mathcal{O}(k) \rightarrow \mathcal{O}(k+1), \quad \partial_k := d^0 - d^1 + \dots + (-1)^{k+1} d^{k+1}$$

satisfies $\partial_{k+1}\partial_k = 0$. Thus we obtain a cochain complex $\{\mathcal{O}, \partial\}$ with *total degree*

$$|x| := \tilde{x} - l \quad \text{for } x \in \mathcal{O}(l)$$

(this agrees with the homological degree in the spectral sequence). We call this the *Hochschild complex associated with \mathcal{O}* .

Define the *normalized Hochschild complex $\tilde{\mathcal{O}}$* by

$$\tilde{\mathcal{O}}(k) := \bigcap_{0 \leq i \leq k-1} \ker\{s^i : \mathcal{O}(k) \rightarrow \mathcal{O}(k-1)\}.$$

The following is a well-known fact.

Lemma 4.1 (see [7, III, Theorem 2.1] for the simplicial version). *The map ∂_k restricts to $\partial_k : \tilde{\mathcal{O}}(k) \rightarrow \tilde{\mathcal{O}}(k+1)$. The inclusion map $\tilde{\mathcal{O}} \rightarrow \mathcal{O}$ is a quasi-isomorphism.*

A Poisson algebra structure on the Hochschild homology $HH(\mathcal{O}) := H_*(\mathcal{O}, \partial)$ was defined in [17]; for $x \in \mathcal{O}(l)$ and $y \in \mathcal{O}(m)$, define two operations

$$\begin{aligned} x \bullet y &:= (-1)^{l\tilde{y}} \mu(x, y) && \in \mathcal{O}(l+m)_{\tilde{x}+\tilde{y}}, \\ [x, y] &:= x\bar{\circ}y - (-1)^{(|x|+1)(|y|+1)} y\bar{\circ}x && \in \mathcal{O}(l+m-1)_{\tilde{x}+\tilde{y}}, \end{aligned}$$

where $\bar{\circ}$ is defined by

$$x\bar{\circ}y := \sum_{1 \leq i \leq l} (-1)^{(m-1)(l-i)+(l-1)\tilde{y}} x \circ_i y,$$

which should be compared with the star-operation $*$ (§3.2).

Theorem 4.2 ([17]). *If \mathcal{O} is a multiplicative operad of graded modules, then $HH(\mathcal{O})$ is a Poisson algebra with respect to \bullet , $[\cdot, \cdot]$ and the degree $|\cdot|$.*

4.2. Connes' boundary operation. Suppose in addition that \mathcal{O} is a cyclic multiplicative operad (see [11, Definition 3.11]); namely, \mathcal{O} is a multiplicative operad with grading-preserving linear maps

$$\tau_k : \mathcal{O}(k) \rightarrow \mathcal{O}(k)$$

satisfying $\tau_k^{k+1} = \text{id}$, $\tau_0(e) = e$, $\tau_2(\mu) = \mu$ and, for $x \in \mathcal{O}(l)$ and $y \in \mathcal{O}(m)$,

$$\tau_{l+m-1}(x \circ_i y) = \begin{cases} \tau_l(x) \circ_{i-1} y & i \geq 2, \\ (-1)^{\tilde{x}\tilde{y}} \tau_m(y) \circ_m \tau_l(x) & i = 1. \end{cases}$$

Lemma 4.3 ([11, Theorem 1.4 (a)]). *Let \mathcal{O} be a cyclic multiplicative operad of graded modules. The collection $\{\tau_k\}_{k \geq 0}$ of maps makes the cosimplicial module \mathcal{O} into a cocyclic module; that is, for $1 \leq i \leq k$, we have*

$$\tau_k d^i = d^{i-1} \tau_{k-1}, \quad \tau_k s^i = s^{i-1} \tau_{k+1}.$$

Define the operation $B_k : \mathcal{O}(k) \rightarrow \mathcal{O}(k-1)$ by

$$(4.1) \quad B_k(x) := (-1)^{\tilde{x}} \sum_{1 \leq i \leq k} (-1)^{i(k-1)} \tau_{k-1}^{-i} s^{k-1} \tau_k(1 - \tau_k)(x).$$

This map is called *Connes' boundary operation* (for non-graded simplicial version, see [9, (2.1.7.1)]). Indeed B is a boundary map:

Lemma 4.4 ([9, §2]). *We have $B_k B_{k+1} = 0$ and $B_{k+1} \partial_k = -\partial_{k-1} B_k$.*

Note that τ_k does not descend to a map on $\tilde{\mathcal{O}}(k)$. But the following holds.

Lemma 4.5 ([9, §2]). *B_k restricts to a map $B_k : \tilde{\mathcal{O}}(k) \rightarrow \tilde{\mathcal{O}}(k-1)$ of the form*

$$(4.2) \quad B_k(x) = (-1)^{\tilde{x}} \sum_{1 \leq i \leq k} (-1)^{i(k-1)} \tau_{k-1}^{-i} \sigma_k(x),$$

where $\sigma_k := s^{k-1} \tau_k$.

The formula (4.1) is equal to (4.2) on $\tilde{\mathcal{O}}(k)$ because $s^{k-1} \tau_k^2 = \tau_{k-1} s^0$ as a consequence of Lemma 4.3.

We have the induced map B_k on Hochschild homology by Lemma 4.4. The main result of this section is the following.

Theorem 4.6. *($HH(\mathcal{O}), \bullet, [\cdot, \cdot], B$) is a BV-algebra with respect to the grading $|\cdot|$.*

This theorem has been already proved for cyclic multiplicative operads of non-graded modules [18], [11, §6]. The proof below is exactly same as that in [11, §6] when the degrees a and b are both even.

Proof. Let $x \in \tilde{\mathcal{O}}(l)_a$, $y \in \tilde{\mathcal{O}}(m)_b$. Define $Z(x, y) \in \tilde{\mathcal{O}}(l+m-1)_{a+b}$ by

$$Z(x, y) := (-1)^{|x||y|+a+b} \sum_{1 \leq j \leq l} (-1)^{j(l+m-1)} \tau_{l+m-1}^{-j} \sigma_{l+m}(y \bullet x)$$

and define $H(x, y) \in \tilde{\mathcal{O}}(l+m-2)_{a+b}$ by $H(x, y) := \sum_{1 \leq j \leq p \leq l-1} H_{j,p}(x, y)$, where

$$H_{j,p}(x, y) := (-1)^{j(l-1)+(m-1)(p+1+l)+lb} \tau_{l+m-2}^{-j} \sigma_{l+m-1}(x \circ_{p-j+1} y).$$

It is not difficult to see that the result follows from the three formulas

$$(4.3) \quad B_{l+m}(x \bullet y) = Z(x, y) + (-1)^{|x||y|} Z(y, x),$$

$$(4.4) \quad (-1)^{|x|} (Z(x, y) - B_l(x) \bullet y) - x \bar{\circ} y \\ = (-1)^b \partial H(x, y) + H(\partial x, y) + (-1)^{l+b+1} H(x, \partial y),$$

$$(4.5) \quad z \bullet w - (-1)^{|z||w|} w \bullet z = (-1)^{|z|} (\partial(z \bar{\circ} w) - (\partial z) \bar{\circ} w - (-1)^{|z|-1} z \bar{\circ} (\partial w)).$$

Indeed, (4.3), (4.4) and (4.5) imply that

$$\begin{aligned} & B_{l+m}(x \bullet y) - (B_l(x) \bullet y + (-1)^{|x|} x \bullet B_m(y) + (-1)^{|x|} [x, y]) \\ &= (-1)^{|x|+b} (\partial H(x, y) + (-1)^b H(\partial x, y) + (-1)^{l+1} H(x, \partial y)) \\ &+ (-1)^{|x||y|+|y|+a} (\partial H(y, x) + (-1)^a H(\partial y, x) + (-1)^{m+1} H(y, \partial x)) \\ &- (-1)^{(|x|+1)|y|} (\partial(B_m(y) \bar{\circ} x) - (\partial B_m(y)) \bar{\circ} x - (-1)^{|y|} B_m(y) \bar{\circ} (\partial x)). \end{aligned}$$

The formula (4.3) follows directly from the definition, and (4.5) is [17, (3.7)]. (4.4) follows from the following formulas, which are proved similarly as in [11, §6]:

- $H(d^0 x + (-1)^{l+1} d^{l+1} x, y) = (-1)^{|x|} Z(x, y) - x \bar{\circ} y,$
- $\sum_{1 \leq j < p \leq l} H_{j,p}((-1)^{p-j} d^{p-j} x, y) = (-1)^{l+b} H(x, d^0 y),$
- $\sum_{1 \leq j \leq p \leq l-1} H_{j,p}((-1)^{p-j+1} d^{p-j+1} x, y) = (-1)^{l+b} H(x, (-1)^{m+1} d^{m+1} y),$
- $\sum_{1 \leq j \leq l} H_{j,l}((-1)^{l-j+1} d^{l-j+1} x, y) = (-1)^{|x|+1} B(x) \bullet y,$
- $\sum_{1 \leq j \leq p \leq l} \sum_{\substack{1 \leq i \leq l \\ i \neq p-j, p-j+1}} H_{j,p}((-1)^i d^i x, y) \\ = (-1)^{b+1} \left(\sum_{1 \leq j \leq p \leq l-1} \sum_{\substack{1 \leq i \leq p-1, \text{ or} \\ p+m \leq i \leq l+m-2}} (-1)^i d^i H(x, y) + d^0 H(x, y) \right. \\ \left. + (-1)^{l+m-1} d^{l+m-1} H(x, y) \right),$
- $\sum_{\substack{1 \leq j \leq p \leq l-1 \\ p \leq i \leq p+m-1}} (-1)^i d^i H_{j,p}(x, y) = (-1)^l \sum_{1 \leq i \leq m} (-1)^i H(x, d^i y). \quad \square$

Corollary 4.7. B_* defines a BV-algebra structure on E^2 -term of the Bousfield homology spectral sequence (which converges to $H_*(\text{EC}(1, B^{n-1}))$) when $n \geq 4$ and descends to a BV-operation on E^∞ -term.

Proof. A cyclic structure on the operad \mathcal{CB}_n of “conformal n -balls” was described in [3]. An easy observation shows that $\tau_* \mu = \mu$ for the operad $H_*(\mathcal{CB}_n) \cong H_*(f\mathcal{C}_n)$, where the multiplication $\mu \in H_0(f\mathcal{C}_n(2)) \cong \mathbb{Z}$ corresponds to $1 \in \mathbb{Z}$. Thus $H_*(f\mathcal{C}_n)$ is a cyclic multiplicative operad of graded modules, and hence $E_{**}^2 = HH_*(H_*(f\mathcal{C}_n))$ admits a BV-algebra structure.

The Bousfield spectral sequence [1] is derived from the double complex $\{C_*(fK_n(*)), d, \partial_*\}$, where fK_n is the framed Kontsevich operad [14] (which is cyclic and multiplicative), C_* is the singular chain complex functor and d is the boundary operator for singular chains. This spectral sequence is a spectral sequence

of Poisson algebras [14, 12]. The map B_* is defined on $C_*(fK_n(*))$ by (4.1) and commutes with both ∂ and d since τ_{k-1} and s^{k-1} are induced by continuous maps defined on $fC(*)$; τ is induced by the cyclic permutation of balls, and s is the forgetting map. Thus B_* commutes with all the differentials d^r on E^r , $r \geq 2$. \square

Conjecture. *At least over rationals, B descends to a map on $H_*(EC(1, B^{n-1}))$ and coincides with Δ discussed in §3.*

ACKNOWLEDGMENT

The author is deeply grateful to Ryan Budney and Paolo Salvatore for informing him about their framed little disks action, to Victor Turchin for answering questions about his Poisson structure on Hochschild homology, and to Takahito Naito for invaluable discussions. The author is partially supported by Grant-in-Aid 228006, 23840015 and 25800038, JSPS.

REFERENCES

- [1] A. Bousfield, *On the homology spectral sequence of a cosimplicial space*, Amer. J. Math. **109** (1987), no. 2, 361–394.
- [2] R. Budney, *Little cubes and long knots*, Topology **46** (2007), 1–27.
- [3] ———, *The operad of framed discs is cyclic*, J. Pure Appl. Algebra **217** (2008), 193–196.
- [4] R. Budney and F. R. Cohen, *On the homology of the space of knots*, Geom. Topol. **13** (2009), 99–139.
- [5] M. Chas and D. Sullivan, *String topology*, math.GT/9911159.
- [6] W. Dwyer and K. Hess, *Long knots and maps between operads*, Geom. Topol. **16** (2012), no. 2, 919–955.
- [7] P. Goerss, J. Jardine, *Simplicial homotopy theory*, Reprint of the 1999 edition, Modern Birkhäuser Classics, Birkhäuser Verlag, Basel, 2009.
- [8] A. Hatcher, *Topological moduli space of knots*, preprint available at <http://www.math.cornell.edu/~hatcher/Papers/knotspaces.pdf>.
- [9] J.-L. Loday, *Cyclic homology*, Grundlehren der Mathematischen Wissenschaften 301, Second Edition, Springer-Verlag, Berlin, 1998.
- [10] J. McClure and J. Smith, *Cosimplicial objects and little n -cubes I*, Amer. J. Math. **126** (2004), no. 5, 1109–1153.
- [11] L. Menichi, *Batalin-Vilkovisky algebras and cyclic cohomology of Hopf algebras*, K-Theory **32** (2004), 231–251.
- [12] K. Sakai, *Poisson structures on the homology of the space of knots*, Geometry and Topology Monographs, Vol. **13** (2008), 463–482.
- [13] ———, *Nontrivalent graph cocycle and cohomology of the long knot space*, Algebr. Geom. Topol. **8** (2008), 1499–1522.
- [14] P. Salvatore, *Knots, operads and double loop spaces*, Int. J. Res. Not. (2006).
- [15] ———, *The topological cyclic Deligne conjecture*, Algebr. Geom. Topol. **9** (2009), 237–264.
- [16] ——— and N. Wahl, *Framed discs operads and Batalin-Vilkovisky algebras*, Quart. J. Math. **54** (2003), 213–231.
- [17] V. Tourtchine, *On the Homology of the Spaces of Long Knots*, in “Advances in Topological Quantum Field Theory”, NATO Sciences series by Kluwer 2005, pp 23–52.
- [18] T. Tradler, *Poincare duality induces a BV-structure on Hochschild cohomology*, Ph. D. Thesis, City University of New York, 2002.

DEPARTMENT OF MATHEMATICAL SCIENCES, SHINSHU UNIVERSITY, 3-1-1 ASAHI, MATSUMOTO, NAGANO 390-8621, JAPAN

E-mail address: ksakai@math.shinshu-u.ac.jp

URL: <http://math.shinshu-u.ac.jp/~ksakai/index.html>