

<Data>

Forest structure in a beech (*Fagus crenata* Blume) stand on a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira, central Japanese snowbelt

Hideyuki IDA

Forest structure in a beech (*Fagus crenata* Blume) stand on a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira, central Japanese snowbelt. Hideyuki IDA (Institute of Nature Education in Shiga Heights, Faculty of Education, Shinshu University, Shigakogen, Yamanouchi-machi, Nagano 381-0401, Japan. E-mail : pida@shinshu-u.ac.jp). *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* 50 : 33-40 (2013).

Nomenclature : Yonekura (supervised by Murata) (2012)

Introduction

Beech (*Fagus crenata* Blume) forests are the most common type of forest vegetation in the cool-temperate zone of Japan. However, in the middle of the 20th century, many beech stands were cut for fuel and pulp production and replaced by coniferous plantations. As a result, the proportion of Japan's forest area covered by natural beech forests has decreased substantially.

Two ecotypes of Japanese beech occur in the country : the Japan Sea type and the Pacific Ocean type. The Japan Sea type occurs in areas with abundant winter snowfall ; it is usually dominated by *F. crenata* in the canopy layer and often forms pure forests. The Pacific Ocean type occurs in areas that do not experience heavy snowfall and is distinguished by the formation of mixed forests with a relatively lower dominance of *F. crenata* (Shimano 2006). The beech forests of Kayanodaira, which was the location for this study, belong to the Japan Sea type.

This paper presents 2012 tree census data and stand structure for a 1-ha permanent plot of beech forest for the Monitoring Sites 1000 Project launched by the Ministry of the Environment of Japan (Ishihara et al. 2011). The 2005–2009 tree census data from the project have already been published as a Data Paper (Ishihara et al. 2011).

Site description

1) Natural environment

The Kayanodaira Heights (at 36°50'N, 138°30'E, and approximately 1400–1700m a.s.l.), belongs to Joshin'etsu-kogen National Park, located near Kijimadaira Village in the northeastern part of Nagano Prefecture. This area is a plateau originating from lava flows. At the study site, the mean annual temperature was 4.9°C and annual precipitation was 1677.5mm (1981–2010) according to Mesh Climate Data 2010 (Japan Meteorological Agency 2012). This region near the Japan Sea is characterized by abundant winter snowfall in which the snow cover usually lasts from November to the following May. The maximum snow depth reaches 3–4m annually.

2) Vegetation history

In most of Kijimadaira Village (350m a.s.l.) up to the Kayanodaira Heights, the natural vegetation is mainly forest dominated by beech. However, the area is currently also covered by various other types of forest such as larch (*Larix kaempferi*) and Japanese cedar (*Cryptomeria japonica*) plantations. Some secondary oak forests dominated by *Quercus serrata* and *Q. crispula*, and secondary birch forests typically dominated by *Betula platyphylla* var. *japonica* and *B. ermanii* also exist. Primary and secondary beech forests are confined

to a limited area above 1400m a.s.l. in the Kayanodaira Heights.

The beech stands surrounding the study plot had previously been utilized for fuelwood production. Many disused coalpits (charcoal kilns) remain within the stands. While the stands are technically secondary forests, they are considered to be old-growth beech stands. The beech-dominated forest is mostly national forest at present and is conserved as the Kayanodaira Natural Recreation Forest.

Fagus crenata-dominated forests in this region belong to the Aucubo-Fagetum crenatae, which is an association attached to the larger Saso kurilensis-Fagion crenatae association (Miyawaki et al. 1994). This association usually grows on thick fertile soils under the mesic conditions of the Japan Sea climate, with its heavy winter snowfall. An undergrowth of dwarf bamboo, including *Sasa kurilensis* and *S. senanensis*, is present. *Sasa*, a monocarpic plant genus, is a typical undergrowth vegetation type in Japanese beech forests and may have a large impact on the regeneration of the forests (Ida and Nakagoshi 1996). The simultaneous death of *Sasa* improves the survival rate of seedlings and a large seedling bank can be produced (Nakashizuka 1988 ; Abe et al. 2001).

Outline of the permanent plot

1) Research history of the plot (Figs. 1, 2)

A permanent plot (named KYD; 36°50'18"N, 138°30'00"E, 1500m a.s.l.; Figs. 1, 2) of 1 ha (100m×100m) in area, horizontally projected on a gentle slope in an old-growth stand dominated by beech (*Fagus crenata*) was originally established in 1998. The plot was enrolled in 2005 as one of the core sites for the Monitoring Sites 1000 Project (Ishihara et al. 2011). The forest floor is characterized by dense dwarf bamboo, *Sasa senanensis*, dominates the plot. Within the plot each year as part of the monitoring project, a tree census is conducted, reproductive organs for beech and fallen litter are observed using litter traps, and ground insects are collected using pitfall traps. In and around the plot, the bird community is also surveyed every year.




Fig. 1. Map of the study site showing the location of a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira. Dashed line shows nature trails.



Fig. 2. The *Fagus crenata* stand in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira in 2012.

In and around the plot, ecological studies have been undertaken since 1995. For example, an approximately 10-ha permanent stand was established to investigate the bird community. Sixty-six bird species have been recorded in the breeding season, with a strong presence of hole-nesting birds (Hotta et al. 2000). Ida et al. (2004) reported the predispersal predation of beech seeds by rodents, as observed using a tree tower set near the plot (Hotta et al. 2001). In 2011 and 2012, the monitoring of carbon dynamics (Hirota et al. 2011) and an investigation of the chemical characteristics of soil organic matter (Iimura et al. 2012, 2013) were conducted in the plot.

2) Vegetation

Fagus crenata is the dominant canopy tree species in the plot, but other arboreal species such as *B. ermanii* and *Aesculus turbinata* sometimes occur in the canopy. *Acer nipponicum*, *A. japonicum*, *A. rufinerve*, *Sorbus commixta*, *Padus grayana*, *Tilia japonica*, *Euonymus macropterus*, *Hydrangea paniculata*, and *Viburnum furcatum* are abundant in the shrub layer. *Dryopteris expansa*, *Thelypteris quelpaertensis*, *Carex multifolia*, *Maianthemum dilatatum*, and *M. japonicum* appear frequently in the herb layer. An undergrowth of *S. senanensis* exists in the stands.

3) Tree census data (Table 1, Figs. 3-6, Supplement)

All living trunks in the plot with a trunk girth at breast height 130cm (GBH) ≥ 15 cm have been tagged with an aluminum plate (Fig. 3a) holding an ID number (A0001-A1164 in 2012; shown in the attached supplement) and identified, and the coordinates (N[x]-W[y] axis; Fig. 3b) for the position of trunk roots have been recorded since 2005. Measurements of GBH (Fig. 4), confirmation of the survival of all tagged trunks, and an investigation of newly recruited trunks have been conducted every autumn (September–November) since 2006. Additionally in 2012, the forest layer to which each tagged trunk belonged was classified as having either a vertically unshaded trunk crown (unshaded area $\geq 30\%$, i.e., shaded area $< 70\%$) or a shaded trunk crown (unshaded area $< 30\%$, i.e., shaded area $\geq 70\%$; Fig. 5). The layers




Fig. 3. (a) An aluminum plate holding the trunk ID number (Racetrack Aluminum Tags, Forestry Supplies Inc.) with a stainless steel nail (#12 \times 50mm) and (b) a red plastic pile (Etapron K-50, Juzen Corp.), driven into the ground at intervals of 10m, showing the coordinates (N[x]-W[y] axis) written with the oil-based marker; Magic Ink, Teranishi Chemical Industry Co., Ltd.) for the position in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira.

were distinguished by their vertical physiognomy as I) the trunk reached the canopy; II) trunk height ≥ 5 m under the canopy or trunk height < 10 m in the fully opened canopy gap; or III) shrub (trunk height < 5 m).

The 2012 measurements of species composition, basal area, density, and characteristics of diameter at breast height 130cm (DBH; GBH/π) of all living trunks (≥ 5 cm DBH) in the 1-ha plot are shown in Table 1. Trunks < 5 cm DBH were omitted in Table 1. In total, 20 woody species were recorded in the plot. A frequency distribution for the DBH of the beech population is shown in Fig. 6. Based on the shape of the distribution, the beech population appears to constantly regenerate.




Fig. 4. Procedure for measuring the girth at breast height 1.3m (GBH) using a steel tape measure (Engineer-pocket 10m: EPK-10, TJM Design Corp.).




Fig. 5. Definitions of the forest layer to which the trunk belonged (I: trunk reached canopy; II: trunk height $\geq 5\text{m}$ under the canopy or trunk height $< 10\text{m}$ in the fully opened canopy gap; III: trunk height $< 5\text{m}$) and whether the trunk crown was vertically shaded (S: shaded area $\geq 70\%$ of the crown, i.e., unshaded area $< 30\%$) or unshaded (U: unshaded area $\geq 30\%$, i.e., shaded area $< 70\%$) in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira.

Table 1. Species composition, basal area, density, and characteristics of diameter at breast height 130cm (DBH) for all living trunks (≥ 5 cm DBH) in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira in 2012.

Species	Family	Basal area (BA)		Density		Density (No./ha) for each layer*1 and crown status*2						Diameter at breast height (DBH : cm)			科名
		m ² /ha	%	No./ha	%	I		II		III		mean \pm SD	Max.	Min.	
						Unshaded	Shaded	Unshaded	Shaded	Unshaded	Shaded				
<i>Fagus crenata</i>	Fagaceae	27.89	83.6	251	25.5	83	3	6	31	18	110	27.4 \pm 25.7	106.7	5.0	ブナ
<i>Betula ermanii</i>	Betulaceae	1.12	3.4	5	0.5	4	-	-	-	-	1	46.7 \pm 25.8	75.7	5.2	ダケカンバ
<i>Aesculus turbinata</i>	Sapindaceae	0.91	2.7	21	2.1	4	-	2	-	6	9	19.4 \pm 13.4	48.8	5.1	トチノキ
<i>Acer nipponicum</i>	Sapindaceae	0.88	2.6	185	18.8	-	-	1	1	101	82	7.4 \pm 2.3	18.8	5.0	ツツカエデ
<i>Chengiotanax sciadophylloides</i>	Araliaceae	0.44	1.3	24	2.4	-	1	3	2	6	12	13.7 \pm 7.0	36.2	6.7	コシアブラ
<i>Acer japonicum</i>	Sapindaceae	0.44	1.3	80	8.1	-	-	1	2	34	43	7.9 \pm 2.8	18.4	5.0	ハウチワカエデ
<i>Hydrangea paniculata</i>	Hydrangeaceae	0.44	1.3	129	13.1	-	-	-	-	60	69	6.5 \pm 1.2	10.6	5.0	ノリウツギ
<i>Viburnum furcatum</i>	Adoxaceae	0.24	0.7	81	8.2	-	-	-	-	24	57	6.0 \pm 0.9	8.8	5.0	オオカメノキ
<i>Sorbus commixta</i>	Rosaceae	0.18	0.5	36	3.7	-	-	-	2	13	23	7.6 \pm 2.2	13.3	5.1	ナナカマド
<i>Rutaceae</i>	Rutaceae	0.16	0.5	10	1.0	1	-	-	-	7	-	12.8 \pm 6.6	24.0	6.4	キハダ
<i>Pheledendron amurense</i>	Rutaceae	0.15	0.5	53	5.4	-	-	-	-	30	23	6.0 \pm 0.8	8.6	5.0	ミズキ
<i>Cornus controversa</i>	Cornaceae	0.09	0.3	1	0.1	-	-	-	1	-	-	34.6	34.6	34.6	アカイダヤ
<i>Acer pictum</i> subsp. <i>maxonii</i>	Sapindaceae	0.09	0.3	21	2.1	-	-	-	-	4	17	7.2 \pm 1.7	10.8	5.1	ウヅミズザクラ
<i>Padus grayana</i>	Rosaceae	0.08	0.2	14	1.4	-	-	-	1	3	10	8.4 \pm 2.2	13.2	5.7	ウリハダカエデ
<i>Acer rufinerve</i>	Sapindaceae	0.07	0.2	19	1.9	-	-	-	-	5	14	6.8 \pm 1.3	9.2	5.0	ヒロハノツリバシ
<i>Etiopymus macropterus</i>	Celastraceae	0.07	0.2	9	0.9	-	-	-	-	4	5	9.4 \pm 3.3	15.5	5.3	シナノキ
<i>Tilia japonica</i>	Malvaceae	0.07	0.2	23	2.3	-	-	-	-	10	13	5.6 \pm 0.6	7.8	5.0	ツノハシバミ
<i>Corylus sieboldiana</i>	Betulaceae	0.06	0.2	16	1.6	-	-	-	-	2	14	5.5 \pm 0.6	6.9	5.0	サワフタギ
<i>Symplocos sawafutagi</i>	Symplocaceae	0.04	0.1	3	0.3	-	-	-	-	1	2	6.7 \pm 0.5	7.3	6.2	ミネカエデ
<i>Acer ischonostii</i>	Sapindaceae	0.01	<0.1	3	0.3	-	-	-	-	3	-	5.6 \pm 0.1	5.7	5.5	ヤマウルシ
<i>Toxicodendron trichocarpum</i>	Anacardiaceae	0.01	<0.1	3	0.3	-	-	-	-	3	-	5.6 \pm 0.1	5.7	5.5	ヤマウルシ
Total		33.37	100.0	984	100.0	92	4	16	37	331	504	12.9 \pm 16.3	106.7	5.0	

*1 I : trunk reached canopy ; II : trunk height ≥ 5 m under the canopy or trunk height < 10 m in the fully opened canopy gap ; III : trunk height < 5 m.*2 Unshaded (unshaded area $\geq 30\%$, i.e., shaded area $< 70\%$) ; Shaded (shaded area $\geq 70\%$ of the crown, i.e., unshaded area $< 30\%$).





Fig. 6. Frequency distribution of *Fagus crenata* living trunks (≥ 5 cm diameter at breast height: DBH, $n=251$) as the most dominant tree in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira in 2012.



Observation area for a subplot

Fig. 7. Definition of the five layers by vertical physiognomy in 400 subplots of $5\text{m} \times 5\text{m}$ in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira. I) canopy (height $\geq 10\text{m}$); II) subcanopy ($\geq 5\text{m}$); III) shrub ($< 5\text{m}$); IV) undergrowth, including woody and herbaceous plants and lianas lower than *Sasa* cover or 1.3m in height without *Sasa* cover; S) *Sasa* cover.

4) Layer structure of the stand (Figs. 7, 8)

The plot was divided into 400 subplots of $5\text{m} \times 5\text{m}$. Each subplot was divided into five layers by vertical physiognomy (Figs. 7, 8): I) canopy (height $\geq 10\text{m}$); II) subcanopy (height $< 10\text{m}$, $\geq 5\text{m}$); III) shrub (height $< 5\text{m}$); IV) undergrowth; or S) *Sasa* cover. The “undergrowth” category included woody and herbaceous plants and lianas lower than *Sasa* cover or 1.3m in height without *Sasa* cover. For each layer, percent coverage was determined by a visual inspection. The field survey was conducted in October 2012, before leaves began to fall.

Acknowledgments


I thank Kazuko Iura and Kaichi Harada for their research assistance. I acknowledge the Hokushin District Forest Office for granting me a permit to survey the study plot. This work is a part of the Ministry of the Environment’s “Monitoring Sites 1000” Project and was partly supported by the Japan Society for the Promotion of Science (JSPS No. 24310004 to M. Hirota).

References


- Abe M, Miguchi H, Nakashizuka T (2001) An interactive effect of simultaneous death of dwarf bamboo, canopy gap, and predatory rodents on beech regeneration. *Oecologia* **127**: 281–286.
- Hirota M, Yashiro Y, Iimura Y, Shizu Y, Ohtuka T, Ida H (2011) Carbon dynamics monitoring in an old-growth beech forest in central Japan I: Practical methods for soil respiration and its spatio-temporal variation. *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* **48**: 9–14 (in Japanese)
- Hotta M, Ezaki Y, Banba T, Imahori R (2000) Breeding bird community of the beech forest of Kayanodaira Heights and its temporal change. *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* **38**: 37–47 (in Japanese with English abstract)
- Hotta M, Ezaki Y, Ida H (2001) Access to tree crowns using a scaffolding system in a beech forest with heavy snowfalls. *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* **39**: 15–17 (in Japanese with English abstract)

- Ida H, Hotta M, Ezaki Y (2004) Predispersal predation by rodents to beechnuts (*Fagus crenata* Blume). *Ecological Research* **19**: 503-509
- Ida H, Nakagoshi N (1996) Gnawing damage by rodents to the seedlings of *Fagus crenata* and *Quercus mongolica* var. *grosseserrata* in a temperate *Sasa* grassland-deciduous forest series in southwestern Japan. *Ecological Research* **11**: 97-103
- Iimura Y, Hirota M, Ida H, Ohtuka T (2012) Chemical characteristics of soil organic matter in an old-growth beech forest in central Japan. *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* **49**: 1-5 (in Japanese)
- Iimura Y, Hirota M, Ida H, Ohtuka T (2013) Comparison of quantity and quality of soil organic carbon between matured and gap areas in an old-growth beech forest. *Journal of Geography* (in press)
- Ishihara MI, Suzuki SN, Nakamura M, Enoki T, Fujiwara A, Hiura T, Homma K, Hoshino D, Hoshizaki K, Ida H, Ishida K, Itoh A, Kaneko T, Kubota K, Kuraji K, Kuramoto S, Makita A, Masaki T, Namikawa K, Niiyama K, Noguchi M, Nomiya H, Ohkubo T, Saito S, Sakai T, Sakimoto M, Sakio H, Shibano H, Sugita H, Suzuki M, Takashima A, Tanaka N, Tashiro N, Tokuchi N, Yoshida T, Yoshida Y (2011) Forest stand structure, composition, and dynamics in 34 sites over Japan. *Ecological Research* **26**: 1007-1008
- Japan Meteorological Agency (2012) Mesh Climate Data 2010
- Miyawaki A, Okuda S, Fujiwara R (1994) *Handbook of Japanese Vegetation*, New revise ed. Shibundo, Tokyo
- Nakashizuka T (1988) Regeneration of beech (*Fagus crenata*) after the simultaneous death of undergrowing dwarf bamboo (*Sasa kurilensis*). *Ecological Research* **3**: 21-35
- Shimano K (2006) Differences in beech (*Fagus crenata*) regeneration between two types of Japanese beech forest and along a snow gradient. *Ecological Research* **21**: 651-663
- Yonekura K (supervised by Murata J) (2012) *An enumeration of the vascular plants of Japan.*, Hokuryuukan, Tokyo


I. Canopy (height ≥10m)




II. Subcanopy (height ≥5m)




III. Shrub (height <5m)



S. Sasa cover



IV. Undergrowth



N (m)




Fig. 8. Percent coverage distribution of each layer in 400 subplots of 5m x 5m in a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira in 2012. I) canopy (height ≥10m); II) subcanopy (height ≥5m); III) shrub (height <5m); IV) undergrowth, including woody and herbaceous plants and lianas lower than *Sasa* cover or 1.3m in height without *Sasa* cover; S) *Sasa* cover.

